arXiv:hep-th/9906092v1 12 Jun 1999

SEMICLASSICAL APPROXIMATION FOR CHERN-SIMONS THEORY AND 3-HYPERBOLIC INVARIANTS

A.A. BYTSENKO, L. VANZO, AND S. ZERBINI

Abstract. The invariant integration method for Chern-Simons theory defined on the compact hyperbolic manifold $\Gamma\backslash\mathbb{H}^3$ is verified in the semiclassical approximation. The semiclassical limit for the partition function is presented. We discuss briefly L^2- analytic torsion and the eta invariant of Atiyah-Patodi-Singer for compact hyperbolic 3-manifolds.

1. Introduction

It is known that topological invariants associated with 3-manifolds can be constructed within the framework of Chern-Simons gauge theory [1]. These values have been specified in terms of the axioms of topological quantum field theory in [2], whereas the equivalent derivation of invariants has also been presented combinatorially in [3, 4], where modular Hopf algebras related to quantum groups have been used. The Witten's (topological) invariants have been explicitly calculated for a number of 3-manifolds and gauge groups [5, 6, 7, 8, 9, 10, 11]. The semiclassical approximation for the Chern-Simons partition function may be expressed by the asymptotics for $k \to \infty$ of Witten's invariant of a 3-manifold M and a gauge group G. Typically this expression is a partition function of quadratic functional. This asymptotics leads to a series of $C^\infty-$ invariants associated with triplets $\{M; F; \xi\}$ with M a smooth homology $3-$ sphere, F a homology class of framings of M, and ξ an acyclic conjugacy class of ortogonal representations of the fundamental group $\pi_1(M)$ [12]. In addition the cohomology $H(M; Ad\,\xi)$ of M with respect to the local system related to $Ad\,\xi$ vanishes.

Date: May, 1999.

We thank Prof. F.L. Williams for useful discussion. First author partially supported by a CNPq grant (Brazil), RFFI grant (Russia) No 98-02-18380-a, and by GRACENAS grant (Russia) No 6-18-1997.

This note is an extension of the two previous papers [13, 14]. Here our aim is to use again the invariant integration method [15, 16] in its simplest form in order to evaluate the semiclassical approximation in the Chern-Simons theory. We do this analysing the partition function related to compact hyperbolic 3-manifolds $\Gamma\backslash\mathbb{H}^3$, where $\mathbb{H}^3$ is the real hyperbolic space and Γ is a co-compact discrete group of isometries (for details see Ref. [17]).

We conclude this section introducing the Witten's invariant defined by the partition function associated with a Chern-Simons gauge theory

$$\mathfrak{W}(k) = \int \mathcal{D}A e^{ikCS(A)}, \qquad k \in \mathbb{Z}. \tag{1.1}$$

The formal integration in (1.1) is one over the gauge fields A in a trivial bundle, i.e. 1-forms on the 3-dimensional manifold X_Γ with values in Lie algebra g of a gauge group $\mathfrak{G}$. The Chern-Simons functional $CS(A)$ can be considered as a function on a space of connections on a trivial principal bundle over a compact oriented 3-manifold X_Γ given by

$$CS(A) = \frac{1}{4\pi}\int_{X_\Gamma} \mathrm{Tr}\left(A\wedge dA + \frac{2}{3}A\wedge A\wedge A\right). \tag{1.2}$$

Let X be a locally symmetric Riemannian manifold with negative sectional curvature. Its universal covering $\widetilde{X}\rightarrow X$ is a Riemannian symmetric space of rank one. The group of orientation preserving isometries $\widetilde{G}$ of $\widetilde{X}$ is a connected semisimple Lie group of real rank one and $\widetilde{X}=\widetilde{G}/\widetilde{K}$, where $\widetilde{K}$ is a maximal compact subgroup of $\widetilde{G}$. The fundamental group of X acts by covering transformations on $\widetilde{X}$ and gives rise to a discrete, co-compact subgroup $\Gamma\subset\widetilde{G}$ such that $X=\Gamma\backslash\widetilde{G}/\widetilde{K}$. Let G be a linear connected finite covering of $\widetilde{G}$, the embedding $\Gamma\hookrightarrow\widetilde{G}$ lifts to an embedding $\Gamma\hookrightarrow G$. Let $K\subset G$ be a maximal compact subgroup of G, then $X_\Gamma=\Gamma\backslash G/K$ is a compact manifold. Let $\mathrm{g}=\mathfrak{k}\oplus\mathfrak{p}$ be a Cartan decomposition of the Lie algebra g of G. Let $\mathfrak{a}\subset\mathfrak{p}$ be a one-dimensional subspace and $J=K\bigcap G_{\mathfrak{a}}$ be the centralizer of $\mathfrak{a}$ in K. Fixing a positive root system of $(\mathrm{g},\mathfrak{a})$ we have the Iwasawa decomposition $\mathrm{g}=\mathfrak{k}\oplus\mathfrak{a}\oplus\mathfrak{n}$. For $G=SO(n,1)$ $(n\in\mathbb{Z}_+)$, $K=SO(n)$, and $J=SO(n-1)$. The corresponding symmetric space of non-compact type is the real hyperbolic space $\mathbb{H}^n$ of sectional curvature -1. Its compact dual space is the unit $n-$ sphere.

Since $CS(A)$ does not contain any metric on X_Γ, the quantity $\mathfrak{W}(k)$ is expected to be metric independent, namely to be a (well-defined) topological invariant of X_Γ. Indeed, this fact has been proved in Refs. [3, 4]. In the limit $k\rightarrow\infty$, the asymptotics of the Witten's invariant (semiclassical approximation of Eq. (1.1)), involves only a partition functions of quadratic functionals [1], namely

$$\sum_{[A_f]} \exp\left(ikCS(A_f)\right) \int \mathcal{D}\omega \exp\left(\frac{ik}{4\pi}\int_{X_\Gamma} \mathrm{Tr}(\omega\wedge d_{A_f}\omega)\right). \tag{1.3}$$

In above equation the sum is taken over representatives A_f for each point $[A_f]$ in the moduli-space of flat gauge fields on X_Γ. In addition the ω are Lie-algebra-valued 1-forms and d_{A_f} is the covariant derivative determined by A_f,

$$d_{A_f}\omega = d\omega + [A_f, \omega]. \tag{1.4}$$

2. Quadratic functional with elliptic resolvent

Let M be a compact oriented Riemannian manifold without boundary, and $n = 2m+1 = \mathrm{dim} M$ is the dimension of the manifold. Let $\chi : \pi_1(M) \longmapsto O(V, \langle\cdot\,,\cdot\rangle_V)$ be a representation of $\pi_1(M)$ on real vectorspace V. The mapping χ determines (on a basis of standard construction in differential geometry) a real flat vectorbundle ξ over M and a flat connection map ∇_p on the space $\Omega^p(M,\xi)$ of differential $p-$ forms on M with values in ξ. One can say that χ determines the space of smooth sections in the vectorbundle $\Lambda^p(TM)^* \otimes \xi$. One can construct from the metric on M and Hermitian structure in ξ a Hermitian structure in $\Lambda(TM)^*\otimes\xi$ and the inner products $\langle\cdot\,,\cdot\rangle_m$ in the space $\Omega^m(M,\xi)$. Thus

$$S_{\mathcal{O}} = \langle \omega, \mathcal{O}\omega\rangle_m, \qquad \mathcal{O} = *\nabla_m, \tag{2.1}$$

where $(*)$ is the Hodge-star map. The map $\mathcal{O}$ is formally self-adjoint with the property $\mathcal{O}^2 = \nabla_m^*\nabla_m$. Suppose that the quadratic functional (2.1) is defined on the space $\mathcal{G} = \mathcal{G}(M,\xi)$ of smooth sections in a real Hermitian vectorbundle ξ over M. There exists a canonical topological elliptic resolvent $R(S_{\mathcal{O}})$, related to the functional (2.1), namely

$$0 \xrightarrow{0} \Omega^0(M,\xi) \xrightarrow{\nabla_0} \ldots \xrightarrow{\nabla_{m-2}} \Omega^{m-1}(M,\xi) \xrightarrow{\nabla_{m-1}} \ker(S_{\mathcal{O}}) \xrightarrow{0} 0. \tag{2.2}$$

Therefore, for the resolvent $R(S_{\mathcal{O}})$, we have $\mathcal{G}_p = \Omega^{m-p}(M,\xi)$ and $H^p(R(S_{\mathcal{O}})) = H^{m-p}(\nabla)$, where $H^p(\nabla) = \ker(\nabla_p)/\Im(\nabla_{p-1})$ are the cohomology space. Note that $S_{\mathcal{O}} \geq 0$ and therefore $\ker(S_{\mathcal{O}}) \equiv \ker(\mathcal{O}) = \ker(\nabla_m)$.

Let us choose an inner product $\langle\cdot\,,\cdot\rangle_{H^p}$ in each space $H^p(R(S_{\mathcal{O}}))$. The partition function of $S_{\mathcal{O}}$ with the resolvent (2.2) can be written in the form (see Refs. [18, 16])

$$\begin{aligned}\mathfrak{W}(k) \equiv \mathfrak{W}(k; R(S_{\mathcal{O}}), \langle\cdot\,,\cdot\rangle_H, \langle\cdot\,,\cdot\rangle) = \left(\frac{\pi}{k}\right)^{\zeta(0,|\mathcal{O}|)/2} & e^{-\frac{i\pi}{4}\eta(0,\mathcal{O})} \\ \times\tau(M,\chi,\langle\cdot\,,\cdot\rangle_H)^{1/2}, & \qquad (2.3)\end{aligned}$$

where $|\mathcal{O}| = \sqrt{\mathcal{O}^2}$ is defined via spectral theory. This is the basic formula one has to evaluate. With regard to the quantity $\tau(M,\chi,\langle\cdot\,,\cdot\rangle_H)$, it is related the Ray-Singer torsion. In fact, if $H^0(\nabla) \neq 0$ and $H^p(\nabla) = 0$ for $p = 1, ..., m$, then the product

$$\tau(M,\chi,\langle\cdot\,,\cdot\rangle_H) = T_{an}^{(2)}(M) \cdot \mathrm{Vol}(M)^{-\mathrm{dim}H^0(\nabla)}, \qquad (2.4)$$

is metric independent [19], i.e. the metric dependence of the Ray-Singer torsion $T_{an}^{(2)}(M)$ factors out as $V(M)^{-\mathrm{dim}_{H^0(\nabla)}}$.

As far as the zeta-function $\zeta(0,|\mathcal{O}|$ is concerned, we recall that there exists $\varepsilon, \delta > 0$ such that for $0 < t < \delta$ the heat kernel expansion for self-adjoint Laplace operators $\mathfrak{L}_p$ is given by

$$\mathrm{Tr}\left(e^{-t\mathfrak{L}_p}\right) = \sum_{0\leq\ell\leq\ell_0} a_\ell(\mathfrak{L}_p)t^{-l} + O(t^\varepsilon). \qquad (2.5)$$

Starting with the formula [18]

$$\zeta(0,\mathfrak{L}_p) = a_0(\mathfrak{L}_p) - \dim(\ker(\mathfrak{L}_p)) = a_0(\mathfrak{L}_p) - \mathrm{dim}H^p(R(S)), \qquad (2.6)$$

one can shown that the zeta function $\zeta(s,|\mathcal{O}|)$ is well-defined and analytic for $\Re(s) > 0$ and can be continued to a meromorphic function on $\mathbb{C}$, regular at $s = 0$ and

$$\zeta(0,|\mathcal{O}|) = \sum_{p=0}(-1)^p(a_0(\mathfrak{L}_p) - \mathrm{dim}H^p(R(S))). \qquad (2.7)$$

Furthermore, the zeta function $\zeta(0,|\mathcal{O}|)$ appearing in the partition function (2.3) can be expressed in terms of the dimensions of the cohomology spaces of $\mathcal{O}$. Indeed, if the dimension of M is odd $(n = 2m+1)$ then for all p $a_0(\mathfrak{L}_p) = 0$, because we are dealing with manifold without boundary. Since $H^p(R(S_{\mathcal{O}})) = H^{m-p}(\nabla)$ (the Poincarè duality) for the resolvent (2.2), it follows that

$$\zeta(0||\mathcal{O}|) = -\sum_{p=0}^{m}(-1)^p\mathrm{dim}H^p(R(S)) = (-1)^{m+1}\sum_{p=0}^{m}(-1)^p\mathrm{dim}H^p(\nabla). \qquad (2.8)$$

Finally, the dependence of the eta invariant $\eta(0|\mathcal{O})$ of Atiyah-Patodi-Singer on the connection map $\mathcal{O}$ can be expressed with the help of the formula for the index of the twisted signature operator for a certain vectorbundle over $M \otimes [0,1]$ (see [20, 21, 22]). Furthermore It can be shown [18] that

$$\eta(s|B) = 2\eta(s|\mathcal{O}), \tag{2.9}$$

where the B are elliptic self-adjoint maps on $\Omega(M,\xi)$ defined on p-forms by

$$B_p = (-i)^{\lambda(p)} \left(*\nabla + (-1)^{p+1}\nabla*\right), \tag{2.10}$$

In this formula $\lambda(p) = (p+1)(p+2)+m+1$ and for the Hodge star-map we have used $*\alpha \wedge \beta = \langle \alpha, \beta\rangle_{vol}$. From the Hodge theory we have

$$\text{dimker}B = \sum_{p=0}^{m} \text{dim}H^p(\nabla). \tag{2.11}$$

3. The case of real compact hyperbolic manifolds

In this section, we shall consider the specific case of a compact hyperbolic 3-manifolds of the form $M = X_\Gamma = \Gamma\backslash\mathbb{H}^3$. If the flat bundle, ξ is acyclic, then for L^2- torsion one gets [23]: $[T_{an}^{(2)}(X_\Gamma)]^2 = \mathfrak{R}_\chi(0)$, where $\mathfrak{R}_\chi(s)$ is the Ruelle function. The function $\mathfrak{R}_\chi(s)$ is an alternating product of more complicate factors, each of which is a Selberg zeta function $Z_p(s;\chi)$. The relation of Ruelle and Selberg functions is:

$$\mathfrak{R}_\chi(s) = \prod_{p=0}^{\text{dim}M-1} Z_p(p+s;\chi)^{(-1)^j}. \tag{3.1}$$

The function $\mathfrak{R}_\chi(s)$ extends meromorphically to the entire complex plane $\mathbb{C}$ [24]. The Ruelle function associated with closed oriented hyperbolic 3-manifold $X_\Gamma = \Gamma\backslash\mathbb{H}^3$ has the form $\mathfrak{R}_\chi(s) = Z_0(s;\chi)Z_2(2+s;\chi)/Z_1(1+s;\chi)$. The analytic torsion for manifold X_Γ has been calculated (in the presence of non-vanishing Betti numbers $b_i \equiv b_i(X_\Gamma) = \text{rank}_{\mathbb{Z}}H_i(X_\Gamma;\mathbb{Z})$) in Refs. [13, 14].

Now we consider the evaluation of eta invariant contribution. With regard to this point, a remarkable formula relating $\eta(s,\mathcal{O})$ to the closed geodesics on X_Γ has been obtained by Millson [25]. More explicitly, Millson has proved the following result for a Selberg type (Shintani) zeta function $\widetilde{Z}(s,\mathcal{O})$.

Let us define a zeta function by the following series, which is absolutely convergent for $\Re(s) > 0$,

$$\text{log}\widetilde{Z}(s,\mathcal{O}) \overset{def}{=} \sum_{[\gamma]\neq 1} \frac{\text{Tr}\tau_\gamma^+ - \text{Tr}\tau_\gamma^-}{|\text{det}(I-P_h(\gamma))|^{1/2}} \frac{e^{-s\ell(\gamma)}}{m(\gamma)}, \tag{3.2}$$

where $[\gamma]$ runs over the nontrivial conjugacy classes in $\Gamma = \pi_1(X_\Gamma)$, $\ell(\gamma)$ is the length of the closed geodesic c_γ (with multiplicity $m(\gamma)$) in the free homotopy class corresponding to $[\gamma]$, $P_h(\gamma)$ is the restriction of the linear Poincaré map $P(\gamma) = d\Phi_1$ at $(c_\gamma, \dot{c}_\gamma) \in TX_\Gamma$ to the directions normal to the geodesic flow Φ_t and $\tau_\gamma^\pm$ is the parallel translation around c_γ on $\Lambda_\gamma^\pm = \pm i$ eigenspace of $\sigma_B(\dot{c}_\gamma)$ (σ_B denoting the principal symbol of $\mathcal{O}$). Then $\widetilde{Z}(z, \mathcal{O})$ admits a meromorphic continuation to the entire complex plane, which in particular is holomorphic at $s = 0$ and

$$\log\widetilde{Z}(0, \mathcal{O}) = \pi i \eta(0, \mathcal{O}). \tag{3.3}$$

Furthermore, it is possible to show that $\widetilde{Z}(s, \mathcal{O})$ satisfies the functional equation

$$\widetilde{Z}(s, \mathcal{O})\widetilde{Z}(-s, \mathcal{O}) = e^{2\pi i \eta(0, \mathcal{O})}. \tag{3.4}$$

Now we have all the ingredients for the evaluation of the partition function (2.3) in terms of L^2- analytic torsion and a Selberg type function. The final result is

$$\mathfrak{W}(k) = \left(\frac{\pi}{k}\right)^{\zeta(0,|\mathcal{O}|)/2} \widetilde{Z}(0, \mathcal{O})^{-1/4} \left[T_{an}^{(2)}(X_\Gamma)\right]^{1/2} [\mathrm{Vol}(\Gamma\backslash G)]^{-\mathrm{dim} H^0(\nabla)/2}, \tag{3.5}$$

where $\zeta(0, \mathcal{O})$ and $\widetilde{Z}(0, \mathcal{O})$ are given by Eqs. (2.8) and (3.2) respectively.

4. Concluding remarks

For a real compact hyperbolic 3-manifold, the formula (3.5) gives the value of the asymptotics of the Chern-Simons-Witten invariant. This is the main result of our paper. The invariant (3.5) involves the L^2- analytic torsion, which can be expressed by means of Selberg zeta functions and a Shintani zeta function $\widetilde{Z}(0, \mathcal{O})$ associated with the eta invariant of Atiyah-Patodi-Singer [20]. Finally we note that the explicit result (3.5) can be very important for investigating the relation between quantum invariants for an oriented 3-manifold, defined with the help of a representation theory of quantum groups [3, 4], and Witten's invariant [1], which is, instead, related to the path integral approach.

References

[1] E. Witten, Commun. Math. Phys. **121** (1989) 351.
[2] G. Moor and N. Seiberg, Commun. Math. Phys. **123** (1989) 77.
[3] N. Reshetikhin and V. Turaev, Commun. Math. Phys. **127** (1990) 1.

[4] N. Reshetikhin and V. Turaev, Invent. Math. **103** (1991) 547.
[5] R. Dijkgraaf and E. Witten, Commun. Math. Phys. **129** (1990) 393.
[6] R. Kirby and P. Melvin, Invent. Math. **105** (1991) 473.
[7] D.S. Freed and R.E. Gompf, Commun. Math. Phys. **141** (1991) 79.
[8] L.C. Jeffrey, Commun. Math. Phys. **147** (1992) 563.
[9] S.K. Rama and S. Sen, Mod. Phys. Lett. **A 8** (1993) 2285.
[10] L. Rozansky, Commun. Math. Phys. **171** (1995) 279.
[11] L. Rozansky, Commun. Math. Phys. **175** (1996) 275.
[12] S. Axelrod and I.M. Singer, J. Diff.Geom. **39** (1994) 173.
[13] A.A. Bytsenko, L. Vanzo and S. Zerbini, Nucl. Phys. B **505** (1997) 641.
[14] A.A. Bytsenko, A.E. Gonçalves and W. da Cruz, Mod. Phys. Lett. A **13** (1998) 2453.
[15] A.S. Schwarz, Commun. Math. Phys. **64** (1979) 233.
[16] D.H. Adams, Phys. Lett. B **417** (1998) 53.
[17] A. A. Bytsenko, G. Cognola, L. Vanzo and S. Zerbini, Phys. Rep. **266** (1996) 1.
[18] D.H. Adams and S. Sen, *"Partition Function of a Quadratic Functional and Semiclassical Approximation for Witten's 3-Manifold Invariant"*, hep-th/9503095.
[19] D. Ray and I. Singer, Adv. Math. **7** (1971) 145.
[20] M.F. Atiyah, V.K. Patodi and I.M. Singer, Math. Proc. Camb. Phil. Soc. **77** (1975) 43.
[21] M.F. Atiyah, V.K. Patodi and I.M. Singer, Math. Proc. Camb. Phil. Soc. **78** (1975) 405.
[22] M.F. Atiyah, V.K. Patodi and I.M. Singer, Math. Proc. Camb. Phil. Soc. **79** (1976) 71.
[23] D. Fried, Invent. Math. **84** (1986) 523.
[24] A. Deitmar, Abh. Math. Sem. **59** (1989) 101.
[25] J.J. Millson, Ann. Math. **108** (1978) 1.

Departamento de Fisica, Universidade Estadual de Londrina, Caixa Postal 6001, Londrina-Parana, Brazil; on leave from Sankt-Petersburg State Technical University *E-mail address:* abyts@fisica.uel.br

Dipartimento di Fisica, Universitá di Trento, and Istituto Nazionale di Fisica Nucleare, Gruppo Collegato di Trento, Italy *E-mail address:* vanzo@science.unitn.it

Dipartimento di Fisica, Universitá di Trento, and Istituto Nazionale di Fisica Nucleare, Gruppo Collegato di Trento, Italy *E-mail address:* zerbini@science.unitn.it